



PubMed

Nucleotide

Protein

Genome

Structure

PopSet

Taxonomy

OMIM

Bc

Search PubMed



for



Limits

Preview/Index

History

Clipboard

Details

About Entrez



Abstract



Sort



Save



Clip Add



Order

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Browser

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

☐ 1: Blood 2002 Feb 1;99(3):1038-43

Related Articles, Links

Full text article at
www.bloodjournal.org

The mechanism of tumor cell clearance by rituximab in vivo in patients with B-cell chronic lymphocytic leukemia: evidence of caspase activation and apoptosis induction.

Byrd JC, Kitada S, Flinn IW, Aron JL, Pearson M, Lucas D, Reed JC.

Division of Hematology-Oncology, The Ohio State University, B302
Starling Loving Hall, 320 W 10th Ave, Columbus, OH 43210, USA. byrd-3@medctr.ohsu.edu

Rituximab is a chimeric monoclonal antibody directed at CD20 with significant activity in non-Hodgkin lymphoma (NHL) and chronic lymphocytic leukemia (CLL). A variety of pathways of tumor cytotoxicity different from cytotoxic chemotherapy have been proposed for this therapeutic antibody including antibody-dependent cellular cytotoxicity and complement-mediated cell lysis. This report describes that a proportion of patients with CLL receiving rituximab treatment have in vivo activation of caspase-9, caspase-3, and poly(ADP-ribose) polymerase (PARP) cleavage in blood leukemia cells immediately following infusion of rituximab. This suggests that apoptosis using a pathway similar to fludarabine and other chemotherapeutic agents is intricately involved in the blood elimination of tumor cells after rituximab treatment. Patients having caspase-3 activation and PARP cleavage in vivo had a significantly lower blood leukemia cell count after treatment as compared to those without caspase activation. Significant down-modulation of the antiapoptotic proteins XIAP and Mcl-1 was also noted, possibly explaining in part how rituximab sensitizes CLL cells to the cytotoxic effect of chemotherapy in vivo. These findings suggest that the therapeutic benefit of antibody-based therapy in vivo for patients with CLL depends in part on induction of apoptosis and provides another area of focus for studying mechanisms of antibody-resistance in neoplastic cells.

Publication Types:

- Clinical Trial
- Multicenter Study

RECEIVED

DEC 16 2002

PMID: 11807010 [PubMed - indexed for MEDLINE]

OFFICE OF PETITIONS

1: Blood 2002 Feb 1;99(3):1038-43

The mechanism of tumor cell clearance by rituximab in vivo in patients with B-cell chronic lymphocytic leukemia: evidence of caspase activation and apoptosis induction.


Byrd JC, Kitada S, Flinn IW, Aron JL, Pearson M, Lucas D, Reed JC.

Division of Hematology-Oncology, The Ohio State University, B302 Starling Loving Hall, 320 W 10th Ave, Columbus, OH 43210, USA. byrd-3@medctr.ohsu.edu

Rituximab is a chimeric monoclonal antibody directed at CD20 with significant activity in non-Hodgkin lymphoma (NHL) and chronic lymphocytic leukemia (CLL). A variety of pathways of tumor cytotoxicity different from cytotoxic chemotherapy have been proposed for this therapeutic antibody including antibody-dependent cellular cytotoxicity and complement-mediated cell lysis. This report describes that a proportion of patients with CLL receiving rituximab treatment have in vivo activation of caspase-9, caspase-3, and poly(ADP-ribose) polymerase (PARP) cleavage in blood leukemia cells immediately following infusion of rituximab. This suggests that apoptosis using a pathway similar to fludarabine and other chemotherapeutic agents is intricately involved in the blood elimination of tumor cells after rituximab treatment. Patients having caspase-3 activation and PARP cleavage in vivo had a significantly lower blood leukemia cell count after treatment as compared to those without caspase activation. Significant down-modulation of the antiapoptotic proteins XIAP and Mcl-1 was also noted, possibly explaining in part how rituximab sensitizes CLL cells to the cytotoxic effect of chemotherapy in vivo. These findings suggest that the therapeutic benefit of antibody-based therapy in vivo for patients with CLL depends in part on induction of apoptosis and provides another area of focus for studying mechanisms of antibody-resistance in neoplastic cells.

Publication Types:
Clinical Trial
Multicenter Study

PMID: 11807010 [PubMed - indexed for MEDLINE]



PubMed Nucleotide Protein Genome Structure PopSet Taxonomy OMIM Bc

Search PubMed for [] Go Clear
Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to File

Text Version

Entrez PubMed
Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services
Journals Database
MeSH Browser
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources
Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

1: Br J Haematol 1998 Jan;100(1):135-41

Related Articles, Links

**Elevated levels of biologically active soluble CD40 ligand in the serum of patients with chronic lymphocytic leukaemia.****Younes A, Snell V, Consoli U, Clodi K, Zhao S, Palmer JL, Thomas EK, Armitage RJ, Andreeff M.**

Department of Hematology, University of Texas M. D. Anderson Cancer Center, Houston 77030, USA.

Chronic lymphocytic leukaemia (CLL) is an indolent lymphoproliferative disorder manifested by low growth fraction and prolonged survival of the malignant cells. The mechanisms that enable CLL cells to live longer and to resist apoptosis remain unclear. Because the malignant CLL cells express CD40 and Fas receptors, which can transduce cell-survival and cell-death signals, we examined the role of CD40 in the growth regulation of CLL cells and its interaction with Fas-mediated and fludarabine-induced apoptosis in vitro. Primary CLL cells underwent spontaneous apoptosis in culture, which was enhanced by exogenous human Fas ligand (FasL) or fludarabine. Exogenous CD40L rescued CLL cells from spontaneous apoptosis in a dose-dependent manner, and caused CLL cells to resist apoptosis induced by FasL or fludarabine. Patients' autologous plasma rescued CLL cells from spontaneous apoptosis, an effect that could be reversed with anti-CD40 ligand (CD40L) antibodies. The levels of soluble CD40 ligand in the sera of 51 CLL patients and 55 healthy donors were determined by enzyme-linked immunosorbent assay. The mean soluble CD40L level in normal donors was 0.29 ng/ml compared to a mean value of 0.80 ng/ml in CLL patients ($P < 0.001$). CD40L up-regulated bcl-X(L) mRNA but not bcl-2 in CLL cells within 3-6 h in culture. Our results demonstrated that serum of patients with CLL contained elevated levels of biologically active soluble CD40L, and that CD40L can prolong survival of CLL cells and mediate their resistance to FasL and fludarabine in vitro.

PMID: 9450802 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to File